Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises




## 2/ Gestion des patients souffrant d'ulcères de jambe.

Le conseil du RSM Venous Forum a créé un document destiné à aider les médecins dans le traitement des patients souffrant d'ulcères de jambe.

## *Management of patients with leg ulcers :*

*The RSM Venous Forum council has created a document that is intended to aid physicians when treating patients with leg ulcers.*



# Management of Patients with Leg Ulcers

## Summary

- Leg ulcers cause great distress to patients and cost the NHS >£1 billion each year.
- The prevalence of leg ulcers is increasing.
- Most patients have an underlying vascular cause for their leg ulcers.
- All patients require specialist assessment and most would benefit from compression and treatment of their veins.
- Despite evidence-based guidelines for referral and treatment, current service provision remains poor.

**Urgent action is needed to ensure that all patients with leg ulceration are offered current best practice**

## The Challenge

- Leg ulcers are non-healing wounds on the lower leg, usually due to a problem with veins (and sometimes arteries).
- Most leg ulcers are caused by chronic venous hypertension.
- Leg ulcers usually take many months to heal.
- Without appropriate care, up to two-thirds of healed ulcers will recur within a year.
- Most patients with leg ulcers are managed in community healthcare settings.
- Primary care data suggest that >50% of patients are not referred and do not receive the care they need.
- Chronic wound care costs between £4.5 - £5.1 billion per year; a third of these wounds are leg ulcers.

## Management Recommendations

1. Every patient with a leg ulcer should have an ankle brachial pressure index (ABPI) assessment ('Doppler') on initial presentation to assess the arterial circulation.

*Rationale:* Doppler assessment of ABPI is a valid and reliable way to detect arterial impairment in the lower limb.

2. All patients with an adequate arterial supply (ABPI>0.9) should be offered effective compression therapy.

*Rationale:* Appropriate compression significantly increases healing of venous ulcers

3. All patients should be referred and have early assessment of their veins using colour duplex ultrasound.

*Rationale:* Duplex examination is the gold-standard method for identifying treatable venous problems.

4. All patients with treatable venous hypertension should be offered minimally invasive endovenous interventions (such as endothermal ablation or foam sclerotherapy).

*Rationale:* Early superficial venous treatment (within 2 weeks) speeds up ulcer healing and halves the risk of ulcer recurrence

## Suggested Patient Pathway

PATIENT PRESENTS WITH LEG ULCER

↓

EARLY ASSESSMENT (INCLUDING ABPI) & APPLY COMPRESSION

↓

EARLY REFERRAL TO VASCULAR SERVICE (prompt assessment including venous duplex)

↓

TREATABLE VENOUS HYPERTENSION? → YES

↓ NO

CONTINUE COMPRESSION & COMMUNITY NURSING CARE

YES → EARLY VENOUS INTERVENTION (WITHIN 2 WEEKS) → CONTINUE COMPRESSION & COMMUNITY NURSING CARE

Academic Department
Direct Line: +44 (0) 20 7290 3918
Direct Fax: +44 (0) 20 7290 2989
Email: venous@rsm.ac.uk
Address: 1 Wimpole Street, London, W1G 0AE
Website: www.rsm.ac.uk Telephone: +44 (0)20 7290 2900
Charity no: 206216 VAT reg no: 524413671

The Royal Society of Medicine

VASCULAR SOCIETY

THE SOCIETY FOR VASCULAR TECHNOLOGY OF GREAT BRITAIN AND IRELAND

@RoySocMed /RoyalSocietyofMedicine Royal Society of Medicine