

FIGURE 3 : Different foam structures obtained by varying the preparation parameters: (a) 3% 20 x C1; (b) 3% 20 x C1; (c) 0.5% 20 x C1; (d) 0.5% 50 x C1; (e) 0.5% 20 x C2; (f) 0.5% 50 x C2.



FIGURE 4 : Histograms of the bubbles volume.



FIGURE 5 : Histograms of the bubbles wall thickness.

Acknowledgements

The Authors wish to thank S. Scozzese for assistance in sample preparation and experimental measurements. Samples quick-freeze and images acquisitions were possible thanks to the support of the Tomolab staff, in particular D. Dreossi and N. Sodini. Images reconstruction and Pore3D computations were conducted with the help of F. Brun and R. Cely Barroso.

References

1. Tessari L., Cavezzi A., Rosso M., Cabrera G. Variables in foam sclerotherapy; literature and experimental data. A.N.Z.J. Phleb. 2008; 11: 83-4.
2. Hamel-Desnos C., Allaert F.A., Benigni J.P., et al. Étude 3/1 mousse de polidocanol 3 %, *versus* 1 % dans la grande veine saphène : premiers résultats. Phlébologie 2005 ; 2 : 175-82.
3. Cabrera J., Cabrera J. Jr. Nuevo metodo de esclerosis en las varices tronculares. Patologia Vasc. 1995; 4: 55-73.
4. Cavezzi A., Tessari L. Foam sclerotherapy and techniques: different gases and methods of preparation, catheter *versus* direct injection. Phlebology 2009; 24: 247-51.
5. Sica M. Guide pratique illustré pour réussir les traitements par sclérothérapie échoguidée à la mousse. Éditions Editcrea, 2009.
6. Wollmann J. The history of sclerosant foams. Dermatol. Surg. 2004; 30: 694-703.
7. Monfreux A. Traitement sclérosants des troncs saphéniens et leur collatérales de gros calibre par la méthode MUS. Phlébologie 1997 ; 50 : 351-3.
8. Sadoun S. Critères pour la fabrication d'une mousse sclérosante de qualité. Phlébologie 2011 ; 64 : 26-31.
9. Ferrara F., Bernbach H.R. Efficacité de la sclérothérapie à la mousse en fonction de l'aiguille utilisée. Phlébologie 2005 ; 58 : 229-34.
10. Morrison N. Foam sclerotherapy, how to improve results and reduce side effects. Phlébologie 2009; 62: 23-34.
11. Coleridge Smith P. Saphenous ablation: sclerosant or sclerofoam? Semin. Vasc. Surg. 2005; 18: 19-24. Review.
12. Bergan J.J., Pascarella L. Severe chronic venous insufficiency: primary treatment with sclerofoam. Semin. Vasc. Surg. 2005; 18: 49-56. Review.
13. Tessari L. Nouvelle technique d'obtention de la scléro-mousse. Phlébologie 2000 ; 53 : 129.
14. Brun F., Mancini L., Kasae P., Favretto S., Dreossi D., Tromba G. Pore3D: A software library for quantitative analysis of porous media. Nuclear Instr. Meth. In Physics Research A 2010; 615: 326-32.
15. Ketcham R.A. Three-dimensional grain fabric measurements using high-resolution X-ray computed tomography. J. Structural Geology 2005; 27: 1217-28.
16. Smith T.H., Schneider E., Odgaard A. Star length distribution: a volume-based concept for the characterization of structural anisotropy. J. Microsc. 1998; 191: 249-57.