Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises

Are graduated compression stockings still essential for VTE prophylaxis in general surgical patients?*

S. L. Schulz[1], B. Strechenesser[2], U. Sachsenger[2], S. Müller[1], D. Andresen[1], U. Beer[2]
Departments of [1]Internal Medicine, [2]General Surgery, Vivantes Medical Center Am Urban, Berlin, Germany

Keywords
Venous thromboembolism, VTE prophylaxis, low molecular weight heparin, graduated compression stockings, pulmonary embolism

Summary
Background: Current guidelines for the prevention of venous thromboembolism in surgical patients continue to recommend the use of graduated compression stockings in addition to low molecular weight heparins (LMWH). Most of the studies documenting a prophylactic benefit from these stockings were carried out before the introduction of LMWH.
Patients, methods: We studied the rate of symptomatic venous thromboembolism (VTE) occurring in patients of a general surgery department during two consecutive time periods of 24 months with different prophylactic regimens. During the first 24 months (Group A), prevention measures consisted of UFH and graduated compression stockings. During the second 24 months (Group B) only LMWH was used.
Results: In Group A there were 3[illegible] patients, with a total of 3[illegible] operations. In Group B there were 2986 patients with a total of 2971 operations. 82.5% of the patients in Group A and 84.3% in Group B received VTE prophylaxis (stockings and UFH in A, LMWH only in B). Symptoms suggesting venous thromboembolism occurred in 45 patients in Group A and in 47 in Group B. Examination of these patients revealed a VTE in 7.4 and 5.8 patients.
Conclusions: The incidence of symptomatic venous thromboembolism in patients of a general surgery department was lower than with previously reported prevention protocols of UFH/LMWH. Graduated compression stockings did not yield additional benefit in our patients.

Phlebologie 2008; 37: 61–67

Schlüsselwörter
Venöse Thromboembolie, VTE Prophylaxe, niedermolekulare Heparine, Kompressionsstrümpfe, Lungenembolie

Zusammenfassung
Hintergrund: Aktuelle Leitlinien zur Venen Thrombose prophylaxe empfehlen noch immer den Einsatz von Thromboseprophylactischen Kompressionsstrümpfen zusätzlich zu niedermolekularen Heparinen (NMH). Fast alle Studien, die einen Nutzen dieser Strümpfe nachweisen sollen, sind vor der Einführung der NMH durchgeführt worden.
Patienten, Methoden: Wir untersuchten die Häufigkeit von symptomatischen Lungenarterienembolien und tiefen venösen Thrombosen bei Patienten einer allgemeinchirurgischen Abteilung über zwei Jahre mit unterschiedlichen Prophylaxe Schemata. Während des ersten 24 Monate (Gruppe A) erhielten die Thromboseprophylaxe mit NMH und Kompressionsstrümpfen, während des zweiten 24 Monate (Gruppe B) wurden nur NMH eingesetzt.
Ergebnisse: In Gruppe A wurden 3181 Patienten bei insgesamt 3850 Operationen, in Gruppe B 2989 Patienten mit insgesamt 2971 Operationen. 82,5% der Patienten in Gruppe A und 84% in Gruppe B erhielten Thromboseprophylaxe (Kompressionsstrümpfe und NMH in A, nur NMH in B). 45 Patienten in Gruppe A und 47 in Gruppe B entwickelten Symptome einer Venenthrombose oder einer Lungenarterienembolie. Bei den weiteren Untersuchungen fand sich in Gruppe A bei 7,4 Prozent und in Gruppe B bei 5,8 Prozent die Diagnose einer VTE.
Schlussfolgerung: Wir zeigen einen deutlich niedrigeren Thromboseprophylaxe mit einer Verwendung von NMH bei sich die Inzidenz symptomatischer Beinvenenthrombosen und Lungenarterienembolien in einer chirurgischen Abteilung sehr gering halten. Der Einsatz von Thromboseprophylaxestrümpfen bringt unter diesen Voraussetzungen keinen zusätzlichen Nutzen.

Sind Kompressionsstrümpfe zur Thromboseprophylaxe in der Allgemeinchirurgie noch erforderlich?

Mots clés
Maladie thrombo-embolique veineuse, prophylaxie, héparines à bas poids moléculaires, bas élastiques à compression dégressive, embolie pulmonaire

Résumé
Historique: Les directives actuelles pour la prévention de la Maladie thrombo-embolique veineuse (MTE) en chirurgie continuent de recommander l'usage de bas à compression d'intensité dégressive associés à un traitement par héparines à bas poids moléculaires (HBPM), en sorte que la plupart des études qui ont montré un bénéfice par la compression élastique, ont été faites avant l'apparition des HBPM sur le marché.
Patients et méthodes: Nous avons étudié le taux de MTE symptomatique survenu dans un département de chirurgie générale au cours de 2 périodes consécutives de 24 mois avec différentes méthodes prophylactiques. Au cours des 24 premiers mois, (Groupe A) on a utilisé des HBPM et des bas à compression dégressive. Au cours des 24 mois suivants, (groupe B) seules des HBPM ont été utilisées.
Résultats: Dans le groupe A 3181 patients ont été traités sur un total de 3850 opérations. Dans le groupe B 2986 patients ont été traités sur un total de 2971 interventions. 82,5% des patients du groupe A et 84% du groupe B ont reçu une prophylaxie de la MTE. Malgré tout, 45 patients ont présenté une symptomatologie suspecte une MTE dans le groupe A et 47 dans le groupe B. Il a été mis en évidence 7 cas de MTE dans le groupe A et 5 dans le groupe B.
Conclusion: The taux de la MTE observé dans un département de chirurgie générale a été inférieur à ceux des protocoles de prévention simple antiprophylaxie de HBPM et de HBPM. Dans ces conditions élastique dégressive ne semblent pas avoir d'intérêt complémentaire chez nos patients.

Les bas à compression élastique dégressive sont-ils essentiels pour la prophylaxie de la maladie thrombo-embolique en chirurgie générale?

* Dieser Aufsatz wurde anlässlich der 49. Jahrestagung der Deutschen Gesellschaft für Phlebologie in Mannheim (Gerhard Communication Award 2007) ausgezeichnet.

Received: November 19, 2007; accepted in revised form: November 21, 2007

Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises

Venous thromboembolism (VTE) continues to account for considerable morbidity, mortality and medical care expense. In the USA the incidence of deep venous thrombosis (DVT) is reported to be approximately 450 000 cases per year and the overall incidence of pulmonary embolism (PE) to be about 595 000 cases per year (8). Numerous studies have indicated that especially surgical and trauma patients are at significant risk for developing venous thromboembolic complications, including pulmonary embolism. Without prophylactic treatment, the risk of VTE (12, 19, 50) is considered to be

- 25–30% in general surgery (8, 13, 46),
- 60% after major trauma (22, 23) and
- up to 45–70% with hip or knee replacements.

Both physical and pharmacological measures aimed at preventing thromboembolic events have been previously evaluated. In earlier studies graduated compression stockings were found to substantially reduce the incidence of VTE in surgical patients (2, 13, 30, 46, 60). During recent years, however, notable developments in pharmacological thromboprophylaxis have been achieved. The introduction of low molecular weight heparins (LMWH), in particular, has resulted in significant advances in treatment modalities so that today they have become the standard for VTE prophylaxis in most countries. The prophylactic properties of LMWHs have yielded equal or even slightly better outcomes than unfractionated heparins. Moreover, they have fewer side effects, do not require laboratory monitoring, and possess pharmacokinetic characteristics which permit their use as a subcutaneously administered, fixed daily dose. LMWHs are reported to reduce the overall incidence of deep vein thrombosis after general surgery by up to 70% (2, 33).

In light of this progress, revaluation of the concomitant use of graduated compression stockings, as it is still recommended in this setting, is warranted.

Patients, methods

Study population and design

The study population consisted of all patients treated in a general surgery department over a period of four years.

With a prospective observational study design, the incidence of symptomatic deep venous thrombosis and pulmonary embolism was investigated with two different VTE prophylaxis regimens. During the

- first 24 months (group A) patients at risk received combined treatment for the prevention of thromboembolism consisting of LMWH and graduated compression stockings,
- second 24 month period (group B) only LMWH was used for prophylaxis.

Treatment

Prophylactic treatment was given to all patients undergoing major surgery (>30 min). The treatment was initiated the day before surgery and was continued for five days thereafter, or until complete remobilization was achieved. The prophylactic measures were also taken in patients with minor surgery and patients who were not actually operated on if they were expected to be immobile for more than 6 hours during the day for more than two days. Patients under the age of 14 did not receive VTE prophylaxis.

The thigh-length compression stockings (T.E.D.® Kendall) in group A were fitted to each patient's legs according to the manufacturer's instructions. The patients were advised to wear them 24 hours a day. Compliance with the wearing of stockings was checked at least twice daily. Patients found without stockings or not wearing them properly on more than two occasions were categorized as noncompliant. Patients with known significant peripheral artery disease (intermittent claudication and/or ABI <0.7) did not receive stockings.

The LMWH employed in both groups was enoxaparin. It was administered according to a simple weight-adjusted protocol. Patients with a body mass index (BMI) of up to 30 kg/m² received 20 mg of enoxaparin and patients with BMI >30 kg/m² received 40 mg – subcutaneously, once daily. Patients in whom the use of LMWH was contraindicated (those with a history of or suspected active HIT II or allergic skin reactions) were treated with 15 mg of hirudin subcutaneously, twice daily.

In group B, no stockings were used. The prophylactic treatment consisted of LMWH or hirudin alone, administered following the same protocol as in group A.

All patients in groups A and B were strongly encouraged to get out of bed and walk as early as possible.

Outcomes

In all patients with clinical symptoms suggesting deep vein thrombosis colour-Doppler ultrasonography of the pelvic and leg veins was done, including compression testing of the leg veins. The leg examination included all veins from the thigh down to the ankle. All sonographic examinations were carried out by two highly-skilled examiners with an ATL / Philips HDI 3000 machine, using a 3–5 mHz curved array scanner for the pelvic veins and a 5–7 mHz phased array scanner for the leg veins. If this examination was temporarily not possible, or if the results were inconclusive, ascending phlebography followed.

Patients with symptoms of pulmonary embolism underwent spiral computed tomography of the lungs. This was judged positive if it revealed distinct filling defects in large vessels. If the diagnosis remained inconclusive after this examination, a ventilation-perfusion lung scan (judged positive only in cases of „high probability" results) and/or a pulmonary angiography was performed.

Results

A total of 6157 patients (Tab. I) were treated in the general surgery department during the study period, with a total of 5861 operations.

Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises

Treatment group characteristics

In group A (treated with stockings and LMWH), there were 3181 patients, 1742 men and 1439 women, with a mean age of 48 years. The mean duration of hospital stay was 7.3 days. 2720 of the A patients had surgery with a total of 3896 operations (Tab. 1, Tab. 2). Four hundred sixty-one patients had no surgery. VTE prophylaxis (Tab. 3) was applied to a total of 2623 (82.5%) of the 3181 patients of period A: to 2422 (89.1%) patients with surgery and to 202 (43.8%) patients without surgery. Fifteen patients in group A could not wear compression stockings due to advanced occlusive artery disease and six others because of extreme obesity. Twenty-eight patients were non-compliant with wearing stockings. Three patients in group A needed to be treated with heparin, instead of LMWH, due to contraindications.

In group B (only LMWH), 2986 patients were treated: 1691 men and 1295 women, with a mean age 49 years. The mean duration of hospital stay was 6.2 days and 2551 patients had a total of 2911 operations. Four hundred thirty-three patients had no surgery. VTE prophylaxis was provided to 2512 (84.0%) of the 2986 patients in period B: to 2309 (90.3%) patients with surgery and to 202 (46.6%) patients without surgery. Two patients had to be treated with heparin, instead of LMWH.

Tab. 1
Patient characteristics, types of surgery

		group A	group B
		[illegible] + stockings	[illegible] only
patients	patient number, n	3181	2986
	men	1742	1691
	women	1439	1295
	age [years]	48	49
	days of stay	7.3	6.2
types of surgery	varicose	202	157
	hernia	149	104
	cholecyst	139	62
	anal-rectal	69	86
	appendicectomy	227	155
	appendicitis	99	266
	urological	370	309
	colon	215	205
	liver	5	4
	gallbladder conventional	44	29
	gallbladder lap	398	386
	pancreatic	3	4
	hysterectomy conventional	357	381
	hysterectomy lap	58	44
	rehabilitation abdominal	82	33
	amputations	130	93
	thoracic	328	302
	thoracic (gynecol. lymphadenectomy)	65	29
	skin (breast, transplantation)	83	81
	mouth, nasal sinuses	167	173
	miscellaneous	35	29
	total	3896	2911

Outcomes (Tab. 4)

Deep vein thrombosis

Thirty-eight patients in group A and 40 in group B developed clinical symptoms suggestive of a deep vein thrombosis which required further investigation. Thirty patients in A and 31 patients in B received color-duplex assisted compression sonography and eight patients in A and nine in B received phlebography. A DVT was confirmed in six patients in group A and in three patients in group B.

Tab. 2
Number of operations per patient

number of operations	group A	group B	p-value
	[illegible] + stockings	[illegible] only	
1	2517	2349	0.659
2	127	124	0.758
3	44	45	0.659
4	18	9	0.598
5	9	9	0.890
6	5	10	0.157
8	1	1	0.302
10	0	1	0.302

Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises

	VTE prophylaxis	group A (2000 + stockings)	group B (2000 only)	p value
patients		2101	2066	
with surgery	–	2120	2053	0.002
	+	1829 (89%)	2016 (98.2%)	0.191
without surgery	–	881	822	0.992
	+	302 (33.8%)	332 (40.7%)	0.209
total	+	2072 (82.7%)	2347 (89%)	0.151

+ = (with/without) VTE prophylaxis

Tab. 1
VTE prophylaxis

	group A (2000 + stockings)	group B (2000 only)	p value
examinations	total/positive	total/positive	
duplex	30/5	37/7	
phlebography	6/1	9/0	
pulm. CT	5/1	6/2	
vent. perf. scan	1/0	0/0	
pulm. angiography	0/0	1/1	
total examinations	40/7	47/10	0.986
patients with VTE	7	5	0.465
event rate	3.22	4.20	0.87

Tab. 2
Examinations and VTE rates

Pulmonary embolism

A pulmonary embolism was suspected in six patients in group A and seven in group B during their hospital stay. 12 of these patients underwent spiral computed tomography of the lungs, one patient had a ventilation-perfusion scan and one needed an additional pulmonary angiogram. The pulmonary embolism was confirmed in one patient in group A and in three in group B.

Characteristics of patients with VTE

All the 12 patients (7 in group A, 5 in group B) with confirmed VTE were receiving prophylaxis at the time the VTE event occurred. The mean duration of hospital stay in these patients was 38 days in group A and 29 days in group B. All patients with PE in both groups had malignant disease.

Discussion

This study, which took place over a period of four years and involved 6167 patients on a general surgery ward, revealed no significant differences in the incidence of symptomatic venous thromboembolism whether graduated compression stockings were used in addition to LMWH prophylaxis, or not.

A review of the literature over the last three decades shows that, despite significant advancements in the prevention of venous thromboembolism, there is still a considerable degree of variation in regard to risk group definition, the number of patients receiving prophylaxis, and prophylactic modalities.

To achieve better standardization in VTE prophylaxis, several consensus conferences have generated guidelines which are aimed at promoting and attaining adequate VTE prophylaxis for the individual patient. Yet, even the latest guidelines concerning the prevention of thromboembolism in surgical patients, namely the updated ACCP guidelines from 2004 (24) and those of the International Consensus Statement dating from 2000 (46), are not completely concordant. In regard to general surgery patients, the former do not recommend a specific prophylaxis in low-risk patients, propose the alternative use of stockings or heparins in moderate- and higher-risk patients, and recommend combination therapy only in very high-risk patients. The latter consider the use of compression stockings in low risk patients, recommend stockings with intermittent pneumatic compression as an alternative to LMWH in moderate risk patients with bleeding tendency and only consider compression stocking as an adjunct to LMWH in high risk patients. In other fields of surgery (for example urologic surgery or gynecologic surgery) there is even less evidence for the necessity of compression stockings. In these fields the recommendations are either extrapolated from general surgery or comments concerning the use of compression stockings are lacking – as is also the case in all recently published studies on VTE prophylaxis in medical patients (36, 41, 52).

There are generally nine studies cited which substantiate significant VTE risk reduction following the use of graduated compression stockings in general surgery (65). All of them were carried out 10 or more years ago (3, 5, 28, 43, 55, 56, 58, 63, 64). The study designs are rather heterogeneous and most of them included a total of less than 100 patients (5, 28, 55, 56, 58). Only two out of the nine studies (3, 28) compared the use of stockings with the absence of prophylactic measures. These found a 50% reduction in the rate of venous thromboembolism in patients who wore stockings. In the seven other studies, a combination therapy was used consisting of either intermittent pneumatic compression (43, 56) or pharmacological methods (dextran, low dose heparin, dihydroergotamine-heparin) together with graduated compression stockings (5, 43, 58, 63, 64). In five of the studies the effectiveness of compression stockings was investigated on only one of the patient's legs; the other leg served as a control (5, 43, 55, 56, 58).

Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises

Several studies in the following years have provided evidence that low dose heparin significantly reduced the incidence of postoperative VTE, and the benefit appeared to increase with the additional use of compression stockings (11, 15, 31, 32, 45). Subsequently, low dose heparin in combination with compression stockings became the most widely adopted measures for VTE prophylaxis in general surgery – a regimen which has been used in many facilities until today (4, 10, 12, 65).

When LMWHs were introduced into VTE prophylaxis in the mid-1990s, they first proved their effectiveness in orthopedic surgery, mainly with hip and knee replacements. Later LMWHs were adopted in general surgery and were combined with the compression stocking prophylaxis already in use. Only four studies can be found which show an advantage to the combination of LMWH with compression stockings over stockings alone – none of them in general surgery (1, 38, 42, 47). In all four studies the group employing compression stockings alone is categorized as a placebo group. However, the incidence of VTE in these placebo groups is actually about the same as in previous studies where no prophylactic measures were used. Only one single, rather small study has compared, as we have, the use of LMWH in combination with compression stockings to LMWH alone for VTE prophylaxis. Kalodiki et al. (34), randomized 78 patients undergoing a total hip replacement into three groups:

- a control group without prophylaxis,
- a group which received enoxaparin once daily,
- a group which received enoxaparin and wore graduated compression stockings (TED) for a period of 8–10 days.

All patients had a preoperative perfusion lung scan and a postoperative perfusion-ventilation scan, as well as bilateral ascending phlebography between days 8 and 12. The DVT rates were 93%, 38% and 25%, respectively. No other study comparing LMWH with and without stockings has been published as of today – neither in general surgery nor in any other medical specialty.

The mechanism of action through which compression stockings prevent thromboembolism remains unclear, though it is thought to be multi-functional with a reduction in venous diameter playing the key role. External compression was shown to reduce the overall cross-sectional area of the lower limb and increase the flow velocity within the veins. Compression also appears to improve the evacuation and coaptation of incompetent and incompletely emptied valvular cusps. Reduction in reflux-related stasis and augmented flow velocity could decrease the risk of thrombus formation by reducing venous wall distension, local contact time, and the concentration of coagulation reactants (2, 14, 57, 61, 66).

Despite their widespread use, there is also no agreement as to the most effective stocking design to achieve the above mentioned effects. Thigh-length TED stockings, which are probably employed in most studies, have intended pressures of 18 mmHg at the ankle, 14 mmHg at the calf, 8 mmHg at the knee, 10 mmHg above the knee, and 8 mmHg at the thigh. It is not known whether these pressures are optimal. Moreover, some authors estimate calf-length stockings to be more effective than thigh-length and some prefer non-elastic ones (2, 9, 26). Stocking design is not always specified in the study protocols and, particularly in some of the newer studies, it is not clear whether or not stockings were used at all (18, 27, 30, 44, 59, 41, 36, 52).

Even those stockings believed to be ideally designed may not achieve the warranted effects. A study measuring pressure gradients in patients wearing prophylaxis stockings showed that 90% of the stockings failed to produce the ideal pressure gradient of 18–14–8 from the ankle to the knee; 54% even produced an inverse gradient (7, 62).

Given the recommended pressure gradients intended to be realized by the design of prophylaxis stockings, it remains obscure why other types of compression stockings, e.g., those recommended for the prevention of post-thrombotic syndrome following a deep vein thrombosis, achieve significantly higher pressure ranges (mostly 40–36–21 mmHg) although they must be worn for much longer time periods. These higher pressure compression stockings fail, however, to prevent DVT recurrence, whereas prophylaxis stockings are claimed to reduce DVT incidence by 50% or more (9, 37).

Finally, prophylaxis stockings were conceived for supine patients (61). However, today most patients are mobilized within only a few hours after surgery. Nevertheless guidelines recommend stockings for 5–7 days (61).

Unlike most other studies, ours did not serially screen all our study patients for deep venous thrombosis or pulmonary embolism. The significance of an asymptomatic VTE detected by screening methods such as ^{125}I fibrinogen marking, plethysmography, contrast phlebography, compression sonography, D-dimer testing, or ventilation-perfusion scans is a matter of debate (16, 35, 40, 54). Conclusive evidence that treatment of asymptomatic patients with VTE detected by these methods results in better outcomes is lacking, not only in regard to non fatal or fatal pulmonary embolism (35), but also in regard to postthrombotic syndrome (25) and pulmonary hypertension (48). Recent study designs have begun to focus more, as we have, on symptomatic thromboembolic events as clinically relevant endpoints (41).

Limitations

We cannot rule out the possibility of diagnostic bias in our study, since the use of stockings cannot be blinded and for organizational reasons we were not able to perform randomization. It is possible that physicians were more likely to order an imaging test for patients who had not received compression stockings than for those who had. Nevertheless, the incidence of symptomatic VTE in our surgical patients was very low with both prophylaxis regimens.

A decline in VTE rates in clinical medicine has been anticipated in recent years (17, 19). Research in this area is lacking, but considering the still doubtless significance of the Virchow triad in regard to VTE pathogenesis, the current prevailing concept of early mobilization is likely to be the most effective measure contributing towards the prevention of thromboembolism in all areas of medicine. New surgical techniques, especially endoscopic and fast track surgery and modern anesthesiologic methods, are

Aucun article ou résumé dans cette revue ne peut être reproduit sous forme d'imprimé, photocopie, microfilm ou par tout autre procédé sans l'autorisation expresse des auteurs et de l'éditeur. Editions Phlébologiques Françaises
No article or abstract in this journal may be reproduced in the form of print, photocopy, microfilm or any other means without the express permission of authors and the editor. Editions Phlébologiques Françaises

also reported to reduce VTE rates (49, 51, 53). Furthermore, complicated risk stratification concepts are increasingly being disregarded in favor of simple and effective protocols like the one we used with our surgical patients (24) and finally, new, even more potent, anti-thrombotic agents are being developed (16, 20, 21, 59).

Conclusion

Our data showed that the rate of symptomatic venous thromboembolism in patients on a general surgery ward is kept low when a concept of early remobilization and a simple, but strictly applied protocol of preventative low molecular weight heparin is implemented. Graduated compression stockings did not yield an additional preventive effect with this regimen.

References

1 Agnelli G, Piovella F, Buoncristiani P, Severi P, Pini M, D'Angelo A, et al. Enoxaparin plus compression stockings compared with compression stockings alone in the prevention of venous thromboembolism after elective neurosurgery. N Engl J Med 1998; 339: 80-85.
2 Agu O, Hamilton G, Baker D. Graduated compression stockings in the prevention of venous thromboembolism. Br J Surg 1999; 86: 992-1004.
3 Allan A, Williams JE, Bolton JF, Le Quesne LP. The use of graduated compression stockings in the prevention of post-operative deep vein thrombosis. Br J Surg 1981; 68: 172-174.
4 Arnoldo BV, Lee TA. Elastic compression stockings for prevention of deep vein thrombosis. Cochrane Database Syst Rev 2000; 3: CD001484.
5 Bergqvist D, Lindblad B. The thromboprophylactic effect of graded compression stockings in combination with dextran. Nr Arch Surg 1984; 149: 1329-1331.
6 Bernstein R, Ram N, Brenner B, Sarig G, Ben-Sela G, Cohen B. Venous sonography for the diagnosis of symptomatic venous thrombosis in patients with cancer undergoing chemotherapy. J Clinicoral Med 2004; 23: 655-658.
7 Bess AJ, Williams S, Cranter A, Blair R, Gregg PJ, Hu AC. Graded compression stockings in elective orthopedic surgery. An assessment of the in vivo performance of commercially available stockings in patients having hip and knee arthroplasty. J Bone Joint Surg Br 2000; 82: 116-119.
8 Bick RL, Kaplan EL. Thromboprophylaxis in surgical patients. Eur J Med Res 2004; 9: 104-111.
9 Brandjes B, Buller HR, Heijboer H, Huisman MV, de Rijk M, Jagt H, et al. Randomized trial of effect of compression stockings in patients with symptomatic proximal vein thrombosis. Lancet 1997; 349: 759-762.
10 Caprini JA, Arcelus J, Sephel LR, Colon PR, Reyna JJ. The use of low molecular weight heparins for the prevention of postoperative venous thromboembolism in general surgery. A survey of practice in the United States. Int Angiol 2002; 21: 76-88.
11 Clagett GP, Reisch JS. Prevention of venous thromboembolism in general surgical patients. Results of meta-analysis. Ann Surg 1988; 208: 227-240.
12 Clagett GP. Prevention of postoperative venous thromboembolism. An update. Am J Surg 1998; 108: 515-522.
13 Cohen GA, Tudor RL, Grice G. Rates of venous thrombosis after general surgery. Combined results of randomised clinical trials. Lancet 1986; 2: 143-146.
14 Coleridge-Smith PD, Harry JD, Scurr JH. Deep vein thrombosis: effect of graduated compression stockings on extension of deep veins of the calf. Br J Surg 1991; 78: 724-726.
15 Collins R, Scrimgeour A, Yusuf S, Peto R. Reduction of fatal pulmonary embolism and venous thrombosis by perioperative administration of subcutaneous heparin. Overview of results of randomized trials in general orthopedic and urologic surgery. N Engl J Med 1988; 318: 1162-1173.
16 Colwell CW, Berkowitz SD, Davidson BL, Lotke PA, Ginsberg JS, Lieberman JR, et al. Comparison of Ximelagatran as a oral direct thrombin inhibitor with enoxaparin for the prevention of venous thromboembolism following total hip replacement. A randomized double-blind study. J Bone Joint Surg Br 2005; 1: 2119-2130.
17 Demé R, Mancy-Jones B. Incidence of thromboembolism after transurethral resection of the prostate (TURP) – a study on TED stocking prophylaxis and literature review. Scand J Urol Suppl 2002; 36: 119-123.
18 Dorfleutner JE, Lindbloom JW, Quinlan DB, Wilkins AR, Crowther MA. Short-duration prophylaxis against venous thromboembolism after total hip or knee replacement. A meta-analysis of prospective studies investigating symptomatic outcomes. Arch Intern Med 2002; 162: 1465-1473.
19 Eichinger S, Kyrle PA. Prevention of deep vein thrombosis in orthopedic surgery. Eur J Med Res 2004; 9: 112-118.
20 Eriksson BI, Ekman SA, Lassen MR, Bergqvist M. Enoxaparin compared with enoxaparin for the prevention of venous thromboembolism after hip fracture surgery. N Engl J Med 2001; 345: 1298-1304.
21 Evans DC, Perry CM, Keefe D. Ximelagatran. Melagatran a review of its use in the prevention of venous thromboembolism in orthopedic surgery. Drugs 2004; 64: 649-679.
22 Geerts WH, Jay RM, Code KI, Chen E, Szalai JP, Saibil EA, et al. A comparison of low-dose heparin with low molecular weight heparin as prophylaxis against venous thromboembolism after major trauma. N Engl J Med 1996; 335: 701-707.
23 Geerts WH, Heit JA, Clagett GP. Prevention of venous thromboembolism. Chest 2001; 119: 132S.
24 Geerts WH, Pineo GF, Heit JA, Bergqvist D, Lassen MR, Colwell CW, et al. Prevention of venous thromboembolism. The Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. Chest 2004; 126: 338s-400s.
25 Ginsberg JS, Brokopp T, Roffey DM, Mai Kassen R, Magner D, Birch J. Post-thrombotic syndrome after hip and knee arthroplasty: a cross-sectional study. Arch Intern Med 2000; 160: 669-672.
26 Hansel MT, Brown DL, Ionescu M, Goldberg PA. Should knee-length replace thigh-length graduated compression stockings in the prevention of deep venous thrombosis? SAR J Surg 2002; 83: 15-16.
27 Hirbom M, Eriksson T, Svartstrom K, Lehmann T, Sarna S, Kaste M. Enoxaparin versus heparin for prevention of deep-vein thrombosis in acute ischaemic stroke: a randomized double-blind study. Acta Neurol Scand 2002; 106: 84-92.
28 Holford CP. Graded compression for preventing deep venous thrombosis. Br Med J 1976; 2: 969-970.
29 Horlander KT, Mannino DM, Leeper KV. Pulmonary embolism mortality in the United States, 1979-1998: an analysis using multiple cause mortality data. Arch Intern Med 2003; 163: 1711-1717.
30 Jeffery PC, Nicolaides AN. Graduated compression stockings in the prevention of postoperative deep venous thrombosis. Br J Surg 1990; 77: 346-361.
31 Kakkar VV, Corrigan T, Spindler J, Howard DB, Flute PT, Clarke RJ, Wessler S, Vine FT. Efficacy of low doses of heparin in prevention of deep vein thrombosis after major surgery. A double blind randomized trial. Lancet 1972; 2: 351-356.
32 Kakkar VV, Corrigan TP, Fossard DP, Sidhu PS, Lewis J. Prevention of fatal postoperative pulmonary embolism by low doses of heparin. Re-appraisal of results of international randomized trial. Lancet 1977; 1: 567-569.
33 Kakkar VV, De Lorenzo F. Prevention of venous thromboembolism in general surgery. Baillieres Clin Haematol 1998; 11: 605-619.
34 Kalodiki EP, Hoppensteadt DA, Nicolaides AN, Sawyer A, Gill K, Regina F, et al. Deep venous thrombosis prophylaxis with low molecular weight heparin and elastic compression patients having total hip replacement. A randomized controlled trial. Int Angiol 1996; 15: 162-169.
35 Kearon C. Noninvasive diagnosis of deep vein thrombosis in postoperative patients. Semin Thromb Hemost 2001; 27: 5-9.
36 Kleber FX, Witt C, Vogel G, Koppenhagen K, Schomacker U, Eberhardt W. A randomized comparison of enoxaparin with unfractionated heparin for the prevention of venous thromboembolism in medical patients with heart failure or severe respiratory disease (The Prince study group). Am Heart J 2003; 145: 614-621.
37 Kolbach DN, Sandbrink MW, Hamulyak K, Neumann HA, Prins MH. Non-pharmacological measures for prevention of post-thrombotic syndrome. Cochrane Database Syst Rev 2002; 1: CD004174.